



SUNCROPS PLUS

FOR PREVENTION OF
STRESS DAMAGE TO PLANTS

UNDER GENERAL GUIDELINES AND PRECAUTIONS

1. Use eye protection and proper gloves and clothing when handling the product. Use only in well-ventilated areas. Avoid contact with eyes or skin. In case of contact with eyes, rinse immediately with plenty of water. If swallowed, seek medical advice immediately and show this container or label.
2. Keep container tightly closed and in a well-ventilated place. Keep only in the original container with the original label.
3. When using do not eat, drink or smoke.
4. Repeat the use of the product in case of rain above 10 mm. Double the dose in case of high extreme environmental temperatures.
5. Wash your hands after manipulation.
6. Do not store or transport together with foods or clothing.
7. To obtain optimal result it is recommended that you a) read the all the instructions, b) leave out the manipulation a some agitation method at mixing and during the application c) leave out the manipulation, the right dose and amount of water.
8. Please, following the label instructions.
9. Suncrops is compatible with most pesticides. Do not mix with clorothalonil or sulfur.
10. Keep out of the reach of children.

If you have any question please contact the technical department. Email : contact@suncrops.cl

www.nutriprove.com

CONTAINS

Aluminum silicate	970 gr/kg
Vegetable acids	3 gr/kg
Processed, refined and enhanced aluminum silicate in the form of wettable powder.	

Suncrops form a reflective barrier film that protects plants against sunburn and heat stress with double action physical and Physiological protection.

RECOMMENDED FOR:

- Hydric stress reduction.
- Solar damage reduction
- Preventing sunburn in plants

The company guarantees the composition and content of this product and also the quality within the due date. The storage and application is the sole responsibility of the user and the company does not assume responsibility for any damage resulting from their improper use, handling and storage.

The seller makes no guarantee, express or implied, on the recommended for use, since its application should be determined in each particular case by a professional. Buyer assumes absolutely all risks of use and management, proceed or not, according to the recommendations on this label. you must require the security seal label, induction seal on drums and inviolable lid on all containers to prevent product adulterated or manipulated by others.

RECOMMENDATIONS FOR USE

Crops	Rate	Comments
Tree crops	<u>Season long</u>	Apply in a water volume according to the tree row volume. Prior to the first heat event.
Pome and stone fruit, nut crops, olives, wine grapes small fruits.	a) Apply 1,25 Kgs/100 lts water every 21 days beginning at petal fall (cap fall) if rainfall or high temperatures occur, re-apply.	On hard wet foliage and grape vines add a non-ionic surfactant. the same at low rate of suncrops and at high water volume
Tropical and subtropical crops avocado, banana, mango, Papaya, Paw paw.	<u>Late season program</u>	Make applications 14 days apart beginning at petal fall (cap fall).
Nursery and ornamental plants.	a) Initial application 2,5kgs/100 lts.	Cherries, stone fruit; apply only early season, up to 2,5 cm fruit diameter or post-harvest.
Citrus	b) Subsequent applications at 15 day intervals at 0,6-1,25 Kgs/100 lts. (0,6 Kgs/100 lts in citrus)	Is recommended to apply with turbo sprayer and "tower". Use the sprayers available depending on the size of the orchard and the size of the trees
Field crops	<u>Season long</u>	Apply the first two applications 14 days apart and prior to the first heat event and should be reduced if crop growth is rapid.
Peppers, Tomatoes potatoes, onions, cucurbits such as cucumber, squash, pumpkin, melon and watermelon.	a) 6-8 Kgs/ha beginning just prior to flowering.	Increase the volume of water used throughout the season based upon plant size.
Vegetable seedlings	b) Re-apply at 15-21 day intervals.	Re-apply at 10 day interval as required to maintain and even coverage on the fruit and foliage.
Pineapple	<u>Sunburn and heat</u>	For transplant shock reduction, apply at 12 kgs/ha to plant in trays.
Cotton	a) Initial application 25 kgs/ha	Cucurbits apply to smooth-skinned cucurbits only
Peanuts	b) Subsequent applications 12 kgs/ha at 15-21 day intervals.	Ensure that fruit is well covered for greatest sunburn reduction. DO NOT TREAT TRANSPLANTS BY DIPPING

Net Contents: 12,5 Kg

Manufactured and Guaranteed by:

NUTRIPROVE S.A.

Distributed by:

